

66330-64

1/14/2014

1/5



UNITED STATES ENVIRONMENTAL PROTECTION AGENCY  
WASHINGTON, D.C. 20460

OFFICE OF CHEMICAL SAFETY  
AND POLLUTION PREVENTION

Dave. G. Bolin, Ph.D.  
Arysta LifeScience North America Corporation  
15401 Weston Parkway, Suite 150  
Cary, North Carolina 27513

JAN 14 2014

Subject: Fluoxastrobin 480 SC Fungicide (Alternate Brand Name Evito 480 SC Fungicide)  
EPA Reg. No. 66330-64  
Your submission dated 9/30/13 and resubmission dated 11/14/13 for a supplemental rice label  
EPA Decision Number: 484477

Dear Dr. Bolin:

The supplemental label referred to above, submitted in connection with registration under the Federal Insecticide, Fungicide and Rodenticide Act (FIFRA), as amended, is acceptable.

One copy of the supplemental label stamped "Accepted" is enclosed for your records. Please submit one copy of the final printed label before the product is released for shipment. If you have any questions, please contact Heather Garvie by phone at: 703-308-0034 or via email at: [garvie.heather@epa.gov](mailto:garvie.heather@epa.gov).

Sincerely,

A handwritten signature in cursive script that reads "Shaja B. Joyner".

Shaja B. Joyner  
Product Manager 20  
Fungicide Branch  
Registration Division

Enclosure: Stamped supplemental label

ACCEPTED  
JAN 14 2014  
Under the Federal Insecticide, Fungicide, and Rodenticide Act, as amended, for the pesticide registered under EPA Reg No. 66330-64

[TEXT IN BRACKETS IS OPTIONAL]



Arysta LifeScience

**SUPPLEMENTAL LABELING**

**GROUP 11 FUNGICIDE**

**EVITO®480 SC Fungicide**

Supplemental Label to Add Directions for Use in Rice.

For use in [Arkansas], [California], [Louisiana], [Mississippi], [Missouri] and [Texas]

**For Agricultural Uses**

This supplemental label expires on February 5, 2016 and must not be used or distributed after this date.

**ACTIVE INGREDIENT:**

Fluoxastrobin: [(1E)-[2-[[6-(2-Chlorophenoxy)-5-fluoro-4-pyrimidinyl]oxy]phenyl] (5,6-dihydro-1,4,2-dioxazin-3-yl) methanone-O-methyloxime].....40.3%

**OTHER INGREDIENTS:**.....59.7%

.....TOTAL: 100.0%

This product contains 3.98 pounds of fluoxastrobin per gallon (478 g per liter)

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**DIRECTIONS FOR USE**

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Before using EVITO 480 SC as permitted according to this supplemental label, read and follow all applicable directions, restrictions, worker protection standard requirements, and precautions on the EPA registered main label for EVITO® 480 SC Fungicide (EPA Reg. No. 66330-64).

This supplemental label must be in the possession of the user at the time of application.

**PRODUCT INFORMATION**

EVITO 480 SC Fungicide is a broad-spectrum fungicide for the control of certain diseases in corn (field, sweet and hybrid seed corn), fruiting vegetables and leaf petiole vegetables, listed low growing berries, peanuts, potato and other tuberous and corm

EVITO 480SC Fungicide  
Supplemental Label  
AD020513

vegetables, rice, soybean, squash /cucumbers subgroup 9B, and wheat. EVITO 480 SC Fungicide works by interfering with respiration in plant-pathogenic fungi, and is a potent inhibitor of spore germination and mycelial growth.

**AERIAL APPLICATION**

**Corn (Field, Sweet and Hybrid Seed Corn), listed Low Growing Berries, Rice [\*], Soybean, Tuberous and Corm Vegetables, and Wheat only**

Aerial applications of EVITO 480 SC Fungicide may be made in spray volumes of 2 or more gallons of water per acre (GPA). Avoid application under conditions when uniform coverage cannot be obtained or when excessive spray drift may occur. Do not apply directly to humans or animals. Aerial applications made to dense canopies may not provide sufficient coverage of lower leaves to provide proper pest control.

**[\*Not approved for use on rice in California.]**

**CHEMIGATION**

**Corn (Field, Sweet and Hybrid Seed Corn), Fruiting Vegetables, Leaf Petiole Vegetables, Tuber and Corm Vegetables, Rice [\*], Soybean, Squash/Cucumber Subgroup 9B[\*], and Wheat only**

Apply EVITO 480 SC Fungicide only through sprinkler type irrigation systems, including center pivot, microjet, wheel lines, lateral move, side roll, or overhead solid set irrigation systems. Do not apply EVITO 480 SC Fungicide through any other type of irrigation system.

**[\*Not approved for use on rice in California.] [\*Not approved for use on squash/cucumber subgroup 9B in California]**

**USE DIRECTIONS FOR SPECIFIC CROPS**

EVITO 480 SC Fungicide provides control or suppression of several important diseases of corn (field, sweet and hybrid seed corn), fruiting vegetables and leaf petiole vegetables, listed low growing berries, peanuts, potato and other tuberous and corm vegetables, rice[\*], soybean, and wheat. When reference is made to disease suppression, suppression can mean either erratic control from good to fair or consistent control at a level below that obtained with the best commercial disease control products.

**[\*Not approved for use on rice in California.]**

## RICE [\*]

Disease Control	Rate to Use	Application Timing and Resistance Management
<b>Sheath/Stem Diseases</b> Sheath blight <i>(Rhizoctonia solani)</i>	3.0 to 4 fl oz/A (0.09 to 0.12 lbs ai/A)	<ul style="list-style-type: none"> <li>For optimal results begin applications preventively and make a second application at a 27 day interval. Use the higher rates and when disease pressure is high</li> </ul>
Aggregate sheath spot <i>(Rhizoctonia oryzae-sativae)</i>  Black sheath rot <i>(Gaeumannomyces graminis va. graminis)</i>  Sheath spot <i>(Rhizoctonia oryzae)</i>  Stem rot <i>(Sclerotium oryzae)</i>	4 to 5.7 fl oz/A (0.12 to 0.18 lbs ai/A)	
<b>Foliar Diseases</b>  Brown leaf spot <i>(Cochliobolus miyabeanus)</i>  Leaf smut <i>(Entyloma oryzae)</i>  Narrow brown leaf spot <i>(Cercospora oryzae)</i>	4 to 5.7 fl oz/A (0.12 to 0.18lbs ai/A)	
<b>Panicle Diseases</b> Kernal smut <i>(Neovossia barclayana)</i> Panicle blast <i>(Pyricularia grisea)</i>	4 to 5.7 fl oz/A (0.12 to 0.18 lbs ai/A)	

[\* Not for Use on Rice in California.]

**RESTRICTIONS AND OTHER INFORMATION:**

- Do not apply more than 11.4 fl oz (0.36 lbs ai) of EVITO 480 SC Fungicide per acre per year.
- There is a maximum number of 2 applications per season, with a minimum retreatment interval of 27 days between applications.
- EVITO 480 SC Fungicide may also be applied through chemigation or by air.

EVITO 480SC Fungicide  
Supplemental Label  
AD020513

- Flood water from treated fields may not be used for irrigation purposes for any food/feed crops.
- Do not apply to rice fields if fields are used for fish/shellfish production.
- Do not apply EVITO 480 SC Fungicide within 28 days of harvest.

**Arysta LifeScience North America, LLC**  
**15401 Weston Parkway, Suite 150**  
**Cary, NC 27513**

EVITO is a registered trademark of Arysta LifeScience North America, LLC  
Arysta LifeScience and the Arysta LifeScience logo are registered trademarks of Arysta LifeScience Corporation



**Arysta LifeScience**